

MedNut Mail

The How, When, Where, Which and Why of pharmacotnutrition

Levodopa +carbidopa and manganese

Y Coleman

6th February 2024

<https://medicationsandnutrition.online/mednutmail/>

Editorial

Is there evidence of a levodopa +carbidopa and manganese interaction?

The total metal content and cellular compartmentalization of several key metals including manganese was examined in those with Parkinsons Disease (PD) (<https://doi.org/10.1039/c7mt00244k>).

The 3 regions selected in this study were because of their responses to Parkinson's disease, and include -

1. Substantia nigra - the primary site of the clinical movement disorder;
2. Fusiform gyrus - associated with visual hallucinations;
3. Occipital cortex – selected as an 'internal control'.

Manganese (Mn) is an essential trace element important in many body functions, including neurotransmitter metabolism and antioxidant functions.

The evidence showed manganese concentrations were significantly higher in both healthy and diseased Substantia nigra than the other selected regions. These findings do not appear to have been subsequently challenged.

Speculatively, are the higher manganese concentrations due to environmental inputs? This is an especially pertinent question given occupational exposure to manganese has been found to cause a form of parkinsonism. Further, chronic exposure to high levels of manganese is associated with a a progressive neurological disorder that manifests similarly to PD, and is commonly known as manganism.

Industrially, manganese is important in steel making and non-ferrous alloys. At a manufacturing level manganese compounds are utilized in a broad range of processes, including the production of batteries, matches, fireworks, porcelain, glass-bonding materials and amethyst glass, pharmaceuticals, animal feeds, ceramics, anti-algal agents, bleaching, disinfectant, fungicides, fertilisers, food additive, food packaging, metal cleaning, nutritional supplement, ore flotation, plant nutrient, preservative for fresh flowers and fruits, purifying natural gas, tanning, textile dyeing and printing, welding, as an oxidising agent, as a catalyst in the chlorination of organic compounds, the production of manganese salts, and the manufacture and drying of paints and varnishes.

The levodopa + carbidopa and manganese interaction is a lack of physiological response to a levodopa + carbidopa-based intervention. One paper suggested a lack

Levodopa + carbidopa and manganese

of response to a levodopa + carbidopa intervention is a likely predictor of manganese-induced parkinsonism.

Clinical Considerations

Not only is it important to know when there is an interaction between a medication and a nutrient, it is also important to know when there is not an interaction.

We also need to be aware of the extent of exposure to environmental inputs such as chronic high manganese intake in order to optimise treatment of those with formal PD, idiopathic PD, and manganism.

Clinical Questions

What actions will you initiate if levodopa + carbidopa is prescribed and there is a poor response?

Will you question whether there has been significant and/or sustained exposure to high levels of manganese?

Conclusions

Is there a levodopa + carbidopa and manganese interaction? To date, there is no evidence of a direct positive or negative interaction.

Disclaimer

The information in this article is provided to support Health Professionals. It is not an exhaustive protocol and Health Professionals are advised that adequate professional supervision is accessed to ensure that Duty of Care obligations with respect to safe administration of medicines is met for each consumer.

Case study

Medical History with Nutritional Aspect

Amputation	<input type="checkbox"/>	Constipation	<input type="checkbox"/>	Dysphagia	<input type="checkbox"/>	MND	<input type="checkbox"/>
Anaemia	<input type="checkbox"/>	CVA	<input type="checkbox"/>	Enteral Feed	<input type="checkbox"/>	MS	<input type="checkbox"/>
Arthritis	<input type="checkbox"/>	CVD	<input type="checkbox"/>	Falls	<input checked="" type="checkbox"/>	Osteoporosis	<input type="checkbox"/>
Cancer	<input type="checkbox"/>	Dementia	<input checked="" type="checkbox"/>	Fracture	<input checked="" type="checkbox"/>	PD	<input type="checkbox"/>
CCF	<input type="checkbox"/>	Dentures	<input checked="" type="checkbox"/>	Frailty	<input checked="" type="checkbox"/>	Pressure Area	<input type="checkbox"/>
Chest Infection	<input type="checkbox"/>	Depression	<input type="checkbox"/>	Gout	<input type="checkbox"/>	Renal	<input type="checkbox"/>
COAD	<input type="checkbox"/>	DM Type 1	<input type="checkbox"/>	Hypertension	<input type="checkbox"/>	Ulcer	<input type="checkbox"/>
Confusion	<input type="checkbox"/>	DM Type 2	<input type="checkbox"/>	Incontinent	<input checked="" type="checkbox"/>	UTI	<input type="checkbox"/>
Food Allergies	dyslipidaemia						
Other:	alcoholism, GORD, deafness, cirrhosis, ascites						

Biochemistry with Pharmaconutrition Consequences

Na:	136	mmol/l	Hb:	141	g/L	Albumin:	25	g/L	BSL:		mmol/l
K:	4.3	mmol/l	Lymph:	1.4		Total Protein:	66	g/L	HbA1C:		
Urea:	2.1	mmol/l	MCV:	97	mmol/l	B12:		pmol/L	INR:		
Creatinine:	0.041	mmol/l	Zn:		umol/l	Folate:		nmol/L	TSH:		mIU/L
Other:	eGFR > 90, Ca 2.02, Ca corr 2.29, phos 1.24, Mg 0.74, Mg corr 0.83										

Medications That May Adversely Affect Nutritional Status

Drug	Vits + Mins	bpp >90%	N/V	C/D	Wt	App	Tst	Thir	Sal	Drlg	d m	Dys	BSL
CAMPRAL		<input type="checkbox"/>	NV	D							<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Fruzemide	(40 mg/day) Ca, Cl, K, Mg, Na,	<input checked="" type="checkbox"/>	NV	CD		↓					<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
Lactulose		<input type="checkbox"/>	NV	D		↓					<input type="checkbox"/>	<input type="checkbox"/>	<input checked="" type="checkbox"/>
LYRICA		<input type="checkbox"/>	NV	CD	↓	↑	<input checked="" type="checkbox"/>		↑		<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input type="checkbox"/>
RE VIA		<input type="checkbox"/>	NV	CD	↕	↓					<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
SPIRACTIN	K, Mg	<input checked="" type="checkbox"/>	NV	D							<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
VITA-D GEL CAPS	(1000 IU/day)	<input type="checkbox"/>									<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
		<input type="checkbox"/>									<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Extra drug:	betavit, vital multivitamin tab												

Levodopa + carbidopa and manganese

Transporter-mediated interactions and nutrients

Transporter	OCT1		OCT2		OCT3		THTR2		OCTN1		OCTN2		OAT1		OAT2	
Nutrients - Substrates	B1, choline		B1, choline		B1, choline, carnitine		B1, B6		carnitine, choline		carnitine, choline		B5, B6, B7, B9		B3, B6, vit C	
Nutrients - Inh									vit D, choline						B3	
DRUG	Sub	Inh	Sub	Inh	Sub	Inh	Sub	Inh	Sub	Inh	Sub	Inh	Sub	Inh	Sub	Inh
Furosemide														Y		Y
Pregabalin									Y							
Spirolactone		Y		Y							Y					
<small>Sub – substrate, Inh – inhibitor, B1 – thiamine, B2 – riboflavin, B3 – niacin, B5 – pantothenic acid, B6 – pyridoxine, B7 – biotin, B9 – folic acid, B12 – cobalamin, NMN – N-methylnicotinamide</small>																

Comments – medication and nutrition impacts only

Data summary

Biochemistry

Recent relevant available biochemistry indicates -

- low albumin – may alter availability of furosemide and spiractin therefore advisable to recheck status;
- low Mg – manifests as confusion, disorientation, personality changes, loss of appetite, depression, muscle cramps, tingling, numbness, hypertension, cardiac dysrhythmia, seizures. Is important in the activation of thiamine, vitamin D, vitamin C, and iodide. Magnesium is an intracellular ion therefore serum levels are unlikely to detect early depletion of status. Cellular magnesium status is unknown whilst magnesium levels within acceptable range however if magnesium levels are low then typically indicates significant cellular depletion and intervention recommended. Both furosemide and spiractin impact magnesium availability therefore advisable to recheck status and if still low then consider a magnesium intervention.

Glycaemia

Currently prescribed 2 medications that alter glycaemia.

Pharmaconutrition

Currently prescribed -

- 6 medications that include nausea, vomiting and diarrhoea as side effects;
- 4 medications that include altered appetite as a side effect;
- 3 medications that include hyponatraemia, hypokalaemia, constipation, dry mouth and thirst as side effects.

Levodopa + carbidopa and manganese

Coffee inhibits vitamin D uptake by osteoblasts (bone builders) by inhibiting their vitamin D receptors, and consequently decreases calcium and zinc absorption.

Currently prescribed vitamin D (1 tab/day). Advisable to check vitamin D levels and if still low then review current vitamin D management strategy.

Chronic use of Coloxyl + senna may promote excessive loss of water and electrolytes, especially potassium, and their regular monitoring recommended.

Frusemide increases urinary excretion of calcium, magnesium, potassium and sodium.

Spiractin impairs zinc status.

Two drugs decrease thiamine availability - being frusemide and spiractin however currently prescribed a high-dose thiamine intervention. Advisable to clarify current thiamine status.

Several of the identified membrane transporters inhibit the absorption and/or organ and cellular uptake of thiamine, choline, carnitine, pyridoxine, pantothenate, biotin, folate, niacin, vitamin C and vitamin D. Consequently, blood test results may be unreliable due to their prevention in entering or exiting relevant organs and cells. Advisable for blood tests to be conducted several hours after administration of relevant prescribed medicines.

Bowel management

Regular aperient prescribed.

Oral + anal PRN interventions prescribed.

No Nurse Initiated interventions administered.

Staff comments

Mr ACT is a small, slender, pleasant man who was lying in bed when I went to speak to him - he told me he is not eating much here, that he is a meat and veg man albeit cutup, that he enjoys desserts and milk drinks.

Mr ACT also referred to his flat and to some of his extended families' difficulties.

Mr ACT's friend advised 10 L fluid were tapped off last week.

Levodopa + carbidopa and manganese

Mr ACT told me the food does have an acceptable taste. However, given his acknowledged unreliability plus the prescription of furosemide, advisable to clarify zinc status.

Observations

Mr ACS was sound asleep when I went to speak to him - he did not waken.

Until recently Mr ACS remained weight stable about 112-113 kg, however recent weights are aberrant and require checking.

Pharmaconutrition comments

Mr ACT's diagnoses include falls - nutritional factors that may be contributing to his falls include -

- potassium - important in muscle function, currently prescribed furosemide therefore advisable to clarify status;
- calcium - more likely to be low if potassium or magnesium low; important in muscle function, currently prescribed furosemide therefore advisable to clarify status;
- vitamin D – associated with muscle weakness and consequently falls; currently prescribed colecalciferol therefore advisable to clarify vitamin D status;
- zinc – can decrease food intake through altered sense of taste and poor appetite, and consequently reduced muscle mass; currently prescribed furosemide therefore advisable to check status;
- magnesium - magnesium is important in vitamin D activation, de novo carnitine production, and muscle function, amongst other functions. Also currently prescribed furosemide which significantly decreases magnesium absorption. Advisable to clarify magnesium status;
- carnitine - carnitine is both absorbed and produced de novo, and is important in a range of muscle functions. Three of the membrane transporters for currently prescribed drugs actually either displace or inhibit carnitine movement therefore advisable to clarify status.

Mr ACT's diagnoses include deafness - nutritional factors to ensure within acceptable ranges include –

- folate - currently prescribed furosemide which inhibits folic acid movement therefore advisable to monitor folate status;

Levodopa + carbidopa and manganese

- vitamin C - inadequate dietary intake associated with deafness; currently prescribed furosemide which inhibits vitamin C movement therefore advisable to monitor vitamin C status;
- zinc - inadequate zinc status has been associated with impaired hearing; currently prescribed furosemide therefore advisable to monitor zinc status;
- thiamine – associated with bilateral hearing loss; currently prescribed furosemide which decreases thiamine availability both directly and indirectly.

As Mr ACT's diagnoses include alcohol abuse and is currently prescribed a high-dose thiamine intervention, advisable to check thiamine status, and if well within, or above, recommended levels then review necessity for its continued prescription. If thiamine status is still low, then advisable to check magnesium status as magnesium is important in thiamine activation. It may also be advisable to administer the thiamine intervention at a significantly different time from furosemide and spironolactone.

What else would you include?


Levodopa + carbidopa and manganese

Please read this as it is important ...

The information in this article is provided to support Health Professionals. It is not an exhaustive protocol and Health Professionals are advised that adequate professional supervision is accessed to ensure that Duty of Care obligations with respect to safe administration of medicines is met for each consumer.


Medications have profoundly and positively changed health outcomes however they do generally come with some nutritional harms. By identifying and addressing the nutritional harms, optimal health outcomes are closer to being achieved.

You may be interested in some of our other products ...



MedNut Mail is our free weekly email that identifies and comments upon some aspect of pharmaconutrition.

[For more information click here.](#)



This resource is for innovative clinicians looking to expand their expertise into the area of drug-nutrition interactions so they can continue to provide their best service to the people in their care.

[For more information click here.](#)